

Contains fluazinam, the active ingredient used in Secure®.



## FUNGICIDE

# DETOUR™ 4 SC

## LEADING THE WAY IN DISEASE RESISTANCE

Let Detour™ 4 SC fungicide lead the way to disease-free turf. With the active ingredient fluazinam, Detour 4 SC attacks turf diseases at full-throttle, protecting your precious greens from unsightly damages that diseases often leave behind. Detour 4 SC demonstrates excellent control of dollar spot that is resistant to several classes of fungicides.

With its multi-site mode of action, Detour 4 SC may be used to prevent or delay the development of resistance to other fungicides making it a perfect addition to your disease management program. Get ahead of the traffic with fall application so your turf can arrive in the spring in pristine condition.

### KEY BENEFITS

- Multi-site contact fungicide
- Aids in resistance management
- Superior control of dollar spot
- Effective on a variety of turf diseases, even those resistant to other fungicides

### KEY USES

- Fairways
- Golf courses
- Greens
- Roughs
- Tees



### PRODUCT NOTES

#### EPA REGISTRATION NUMBER

91234-34

#### ACTIVE INGREDIENT

Fluazinam 40%

#### FORMULATION

Suspension concentrate

#### FRAC NUMBER

29

#### SIGNAL WORD

Caution

#### PACKAGE SIZE

2 x 2.5 gal

#### RESTRICTED USE

No

Refer to label for specific use restrictions.



[www.AtticusLLC.com](http://www.AtticusLLC.com)

### APPLICATION INFORMATION

Detour™ 4 SC may be applied to all types of spray equipment normally used for ground applications on golf courses. For optimal disease control, apply with sufficient water to ensure uniform coverage. Apply after mowing or allow treated area to dry completely before mowing or for foliar disease, allow treated area to dry before irrigation. Atticus recommends using Detour 4 SC in conjunction with turf management practices that promote good plant health and optimal disease control.

USE RATES			
DISEASES CONTROLLED PATHOGEN(S)	APPLICATION INTERVALS (DAYS)	RATE FL OZ DETOUR 4 SC PER 1000 SQ FT	INSTRUCTIONS FOR USE ON FAIRWAYS, GOLF COURSES, GREENS, FAIRWAYS, TEES AND ROUGHS
<b>Algal scum</b> <i>(filamentous blue-green algae/cyanobacteria)</i>	14	0.5	This product should be tank-mixed with another fungicide labeled for control of this disease.
<b>Anthracnose</b> <i>Colletotrichum cereal</i>	14	0.5	Applications should be made prior to infection. Alternate or tank-mix this product with another fungicide for anthracnose disease prevention. This product should be tank-mixed with the labeled rate of another fungicide labeled for control of this disease.
<b>Brown blight, leaf spot, melting-out</b> <i>Drechslera spp.</i> <i>(including D. poae, D. siccans)</i>	14	0.5	Apply when conditions become favorable for disease. For broad-spectrum disease control, tank-mix with the labeled rate of another fungicide labeled for control of these diseases.
<b>Brown patch, large patch</b> <i>Rhizoctonia solani</i>	14	0.5	Begin applications in May or June before disease is present for brown patch. To control large patch, apply in Fall when temperature drops below 70°F. For broad-spectrum control or for control greater than 14 days, tank-mix with the labeled rate of another fungicide labeled for control of these diseases.
<b>Dollar spot</b> <i>Sclerotinia homeocarpa</i> <i>Lanzia spp.</i> <i>Moellerodiscus spp.</i>	14	0.5	Apply as a preventive disease program, starting when conditions become favorable for disease. For broad-spectrum disease control or for control greater than 14 days, tank-mix with the labeled rate of another fungicide labeled for control of this disease.
<b>Gray Snow Mold</b> <i>Typhula spp.</i> <b>Pink Snow Mold</b> <i>Microdochium nivale</i>	Late Fall	0.5	Apply one to two applications in the late Fall before snow cover. Do not apply on top of the snow. For optimum disease control, tank-mix with the labeled rate of another fungicide labeled for control of these diseases.
<b>Pink patch</b> <i>Microdochium nivale</i>	14	0.5	Apply in the Fall or early Spring when conditions are favorable for disease development. For optimum disease control, tank-mix with the labeled rate of another fungicide labeled for control of this disease.
<b>Red Thread</b> <i>Laetisaria fuciformis</i>	14	0.5	Apply when conditions become favorable for this disease.
<b>Stem rust</b> <i>Puccinia graminis</i> <i>var. graminicola</i>	14	0.5	Apply on a preventive schedule when disease conditions are favorable (cool and cloudy weather).

Refer to product label for complete application and mixing instructions.

17319 ©2017 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Detour™ 4 SC is not manufactured or distributed by Syngenta, seller of Secure®. Secure® is a registered trademark of Ishihara Sangyo Kaisha. Detour™ is a trademark of Atticus, LLC.

### KEY DISEASES

- Anthracnose
- Brown patch
- Dollar spot
- Leaf spot
- Pink patch
- Pink snow mold

(Refer to product label for complete list)