

Contains fluroxypyr and clopyralid, the active ingredients used in WideMatch®.



SELECTIVE HERBICIDE



UNMATCHED CONTROL OF BROADLEAF WEEDS

WhipLash™ contains fluroxypyr and clopyralid, two of the most powerful herbicides in one easy-to-use formulation. WhipLash is highly effective at controlling some of the toughest weeds that plague growers today. It controls over 60 broadleaf weeds including herbicide-resistant kochia. WhipLash is highly compatible with grass herbicides, resulting in reduced time spent in the field. Plus, the convenient pre-mix formulation offers a flexible application window with relatively low-use rates.

KEY BENEFITS

- Outstanding control of a variety of tough, broadleaf weeds
- Effective against ALS and dicamba resistant kochia
- Two powerful herbicides in one convenient formulation
- Flexible application timing and tank-mix compatible

KEY USES

- Barley
- Corn
- CRP
- Non-cropland
- Oats
- Wheat

PRODUCT NOTES

EPA REGISTRATION NUMBER

91234-46

ACTIVE INGREDIENT

Fluroxypyr 12.3%

Clopyralid 11.3%

FORMULATION

Emulsifiable concentrate

HRAC NUMBER

4

SIGNAL WORD

Caution

PACKAGE SIZE

2 x 2.5 gal

265 gal

RESTRICTED USE

No

Refer to label for specific use restrictions.



Relevant. Simple. Reliable.



APPLICATION INFORMATION

For optimal results, apply WhipLash™ to actively growing weeds during ideal weather conditions. Herbicidal activity is influenced by weather conditions. The ideal temperature range for herbicidal activity is 55°F to 75°F. WhipLash is rainfast in as little as six hours after application.

WHEAT (INCLUDING DURUM), BARLEY AND OATS APPLICATION TIMING

Apply WhipLash as a broadcast post-emergence treatment to actively growing wheat, barley or oats, from the three leaf crop growth stage up to and including flag leaf emergence. Apply when weeds are actively growing, but before weeds are four inches tall or vining. To obtain season-long control of perennial weeds such as Canada Thistle, apply when the majority of the basal leaves have emerged from the soil up to bud stage. For suppression of volunteer potatoes, apply before potato plants are six inches tall.

BROADCAST APPLICATION RATES

WEED SIZE OR SPECIES ¹	APPLICATION RATE (PT/ACRE)
Susceptible broadleaf weed seedlings less than 4 inches tall ²	1.0
Susceptible broadleaf weed seedlings less than 8 inches tall or vining; dicamba tolerant kochia biotypes	1.33
Volunteer potatoes, mayweed chamomile (dog fennel), pineappleweed	1.33

¹ See "Weeds Controlled or Suppressed" section on product label for a complete listing of weeds controlled or suppressed.
² A rate of one pint per acre will provide satisfactory control of kochia seedlings less than four inches tall (including ALS resistant biotypes). However, when conditions for control are less favorable, such as under drought or cool temperatures, a rate of 1.33 pints per acre will provide more consistent control of kochia seedlings one to four inches tall. Control of small kochia will be more consistent if kochia is at least one inch tall. A rate of 1.33 pints per acre should be used for optimal control of dicamba tolerant kochia populations (see "Management of Kochia Biotypes" in the "Broadleaf Weeds Controlled" section on product label).

PRECAUTIONS

- Only weeds emerged at the time of application will be controlled.
- Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.

RESTRICTIONS

- Do not apply more than 1.33 pint of this product per acre per growing season.
- Do not allow livestock to graze treated areas or harvest treated forage within seven days of application.
- Pre-harvest Interval: Do not apply closer than 14 days before cutting of hay or 40 days before harvesting of grain and straw.
- Do not use if cereal crop is underseeded with a legume.
- When tank-mixing, do not exceed label application rates.

Refer to product label for complete application and mixing instructions.

©2018 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. WhipLash™ is a trademark of Atticus LLC. WhipLash™ is not manufactured, or distributed by Dow Agrosciences LLC, seller of WideMatch®. WideMatch® is a registered trademark of The Dow Agroscience LLC.

KEY WEEDS

- Bedstraw
- Buckwheat
- Chickweed
- Clover
- Cocklebur
- Coffeeweed
- Flax
- Horseweed
- Jimsonweed
- Kochia
- Locoweed
- Mallow
- Morningglory
- Nightshade
- Purslane
- Ragweed
- Sunflower
- Thistle
- Velvetleaf

SUPPRESSION ONLY:

- Bindweed
- Horsetail, field
- Knapweed
- Knotweed
- Ladysthumb
- Mallow
- Smartweed
- Volunteer potato

(Refer to product label for complete list)

